



# Plagiarism Checker X Originality Report

**Similarity Found: 23%**

Date: Wednesday, January 23, 2019

Statistics: 607 words Plagiarized / 2618 Total words

Remarks: Medium Plagiarism Detected - Your Document needs Selective Improvement.

---

J. Trop. Plant Prot. 1 (1): 46-51 (2015) 46 Original Article Leaf Extract of *Mirabilis jalapa* L. Induced Defense of Tomato Plant (*Lycopersicon esculentum* Mill.) **Against Cucumber Mosaic Virus (CMV) Infection** Dian Eka KUSUMAWATI, Tutung HADIASTONO, and Luqman Qurata AINI \* Plant Protection Department, Faculty of Agriculture, University of Brawijaya Jl.Veteran, Malang 65145, East Java, Indonesia (Received 07 April 2015; Accepted 1 Mei 2015) Abstract **Infection of Cucumber Mosaic Virus (CMV) could reduce the tomato production up to 50% depending on the age of the plant during infection and tomato varieties.**

Until now the control efforts against CMV infection have been made, but no consistent effective method was available. Recently, new approach by increasing the plant health to defense against plant disease was increased. It can be stimulated by inoculation of microbes associated with roots or by application of plant extract such as *Mirabilis jalapa*.

In this study, the effect of *M. jalapa* leaf extract in different concentration to develop control against CMV infection in tomato plants was conducted. The results showed that the application of *M. jalapa* leaf extract could inhibit the development of CMV disease in tomato plants. The higher the concentration of *M.*

*jalapa* leaf extract applied on tomato plant, the lower disease incidence as well as the longer incubation period of CMV disease in tomato plants. The applications of *M. jalapa* leaf extract elevated the activity of peroxidase and catalase enzymes. Thus, the inhibition of CMV disease development in tomato plant by application of *M.*

*jalapa* leaf extract may related to the induction of peroxidase and catalase activity in

tomato plants. Keywords : M. jalapa, CMV, plant defense, leaf extract, peroxidase, catalase \* To whom correspondence should be addressed at: E-mail: luqman.fp@ub.ac.id INTRODUCTION Tomato is a horticultural commodity which has high economic value.

One important obstacle on the cultivation of tomato plants is the disease caused by plant pathogenic virus. There are 18 types of viruses have been reported could infect tomato plants. Among them Cucumber Mosaic Virus (CMV) is the most harmful one (Crescenzi et al., 1993). CMV infection on tomato caused necrotic spots on fruit, leaf malformation, and in severe condition could lead to stunting and plant death (Crescenzi et al.,

1993; Jorda et al., 1992). On several tomato production sites, the CMV infection caused substantial losses of production (Alonso-Prados et al., 1997). In Indonesia, the virus that commonly infect tomato plants is Cucumber Mosaic Virus (CMV). Infection of CMV could reduce the tomato production up to 50% depending on the age of the plant during infection and tomato varieties.

Until now the control efforts against CMV infection have been made, but no consistent effective method was available (Gallitelli, 1998; Jorda et al., 1992; Summers et al., 1995). CMV disease control efforts have been carried out, either by controlling the vector carrying the virus, as well as using a physical barrier, but the success of these two methods have not been consistent nor effective. Recently, new approach by increasing the plant health to defense against plant disease was increased.

It can be stimulated by inoculation of microbes associated with plant roots or by application of plant extract such as *Mirabilis jalapa* (Hersanti 2004). The ability of plant extracts in inducing plant defense against disease due to the presence of active compounds that could elevate plant defense related genes in the plant (Murphy et al., 2001). *J. Trop. Plant Prot.* 1 (1): 46-51 (2015) 47 In this study the effect of M.

*jalapa* leaf extract in different concentration was conducted to develop control against CMV infection in tomato plants MATERIALS AND METHODS Preparation of *M. jalapa* leaf extract *M. jalapa* leaf extract was prepared by homogenizing the *M. jalapa* leaves in mortar pestle, and the phosphate buffer was then added. The *M. jalapa* leaf extracts at the concentration of 12,5%, 25%, 50%, and 100% were then prepared by adjustment of additional phosphate buffer.

Pot Experiment in Screen House The experiment was conducted using completely randomized design (CRD) with 5 treatments and 5 replications. The composition of the

treatments was as follows: A0 = Without M. jalapa leaf extract (aquadest) A1 = 12.5% concentration of M. jalapa leaf extract A2 = 25% concentration of M. jalapa leaf extract A3 = 50% concentration of M. jalapa leaf extract A4 = 100% concentration of M.

jalapa leaf extract The data were analyzed using the ANOVA test with a significance level of 5%, and then the significant data were further analyzed by HSD test with a significance level of 5%. M. jalapa leaf extract or aquadest (as control) was applied on two leaves above the cotyledons of 4 weeks old tomato plants. After 30 minutes the leaves were rinsed with water.

Inoculation of CMV was done 24 hours after application of the extract by applying the sap of CMV infected leaves on the third and fourth leaves above the cotyledons. The parameters measured were the symptoms shown on tomato test plant, incubation period, and the disease intensity. The disease intensity was calculated as described by Abadi (2003) below.

$I = \frac{\sum (nxv)}{N \times Z} \times 100\%$   
N = number of leaves with each scale of disease severity  
v = severity scale of each leaf  
N = number of leaves per plant observed  
Z = the highest scale of disease severity level  
The scale of disease severity was as follow: 0 = healthy leaf 1 = mosaic lesion at = 25% on leaf 2 = mosaic lesion on leaf at > 25% - = 50% with blister 3 = mosaic lesion on leaf at > 50%, with blistering and malformation 4 = malformation, leaf scald and dwarf  
The activity of catalase and peroxidase enzymes  
The activity of peroxidase and catalase enzymes were measured using a method described by Reddy et al. (1995) and Luck (1974). The leaf samples of tomato plant were collected at 0, 7, and 14 days after inoculation of CMV.

**RESULTS AND DISCUSSION** Symptoms and Incubation Period of CMV on Tomato Plant  
CMV is a systemic viral disease which the symptoms spread to all parts of the plant. In this study the CMV symptom showed initially in young emergence leaves of tomato plants. The leaves infected by CMV showed mosaic symptom and on the later stage, leaves showed malformation to be wrinkle and smaller leaves.

Further development of CMV symptom on severe condition showed yellowing, smaller leaves and stunting on all parts of tomato plant. These conditions were different with non infected tomato plant which showed healthy and green leaves of tomato plants (Figure 1). Lecoq et al. (1998) reported that plant leaves affected by CMV showed a mosaic, necrosis, and malformation.

Further development showed that leaf size tends to shrink, leaves turn yellow and slightly thickened, and the fruits were changed in color and shape. The observation of

the incubation period of CMV symptoms on tomato plants was listed J. Trop. Plant Prot. 1 (1): 46-51 (2015) 48 in Table 1. Based on data shown in Table 1, the treatment of M. jalapa leaf extract significantly affected the period of incubation of CMV on tomato plant.

Tomato plants without application of M. jalapa leaf extract showed the faster incubation period of CMV disease when compared to that of tomato plants treated with leaf extract of M. jalapa. The data also showed that the higher concentration of M. jalapa leaf extract application, the longer the incubation period of CMV disease.

The longest incubation period of CMV was shown in tomato plants treated with 50% and 100% of M. jalapa leaf extract. The result above showed that the treatment of M. jalapa leaf extract could delay the symptom of CMV on tomato plants. This result was similar to the result of Hersanti (2004) which has reported that leaf extract of M. jalapa was able to delay the incubation period of CMV.

(a) (b) (c) (d) Figure 1. Symptoms of CMV infection in tomato plant: (a) mosaic on young leaves; (b) malformation; (c) leaf curling and stunted; (d) healthy leaf Table 1. Incubation Period of CMV Symptom on Tomato Plants Treatment Incubation period (days) No Treatment of M. jalapa leaf extract 8.8 a 12.5% of M.

jalapa leaf extract 11.8 b 25% of M. jalapa leaf extract 13.8 c 50% of M. jalapa leaf extract 20 d 100% of M. jalapa leaf extract 20.4 d Description: The number followed by the same letter in the same column indicates no significant difference based on HSD test (5%). Table 2. The Disease Intensity of CMV On Tomato Plants Treatment Disease Intensity of CMV On Tomato Plants No Treatment of M. jalapa leaf extract 50.158 d 12.5% of M.

jalapa leaf extract 45.712 c 25% of M. jalapa leaf extract 41 226 b 50% of M. jalapa leaf extract 33 948 a 100% of M. jalapa leaf extract 31 646 a Description: The number followed by the same letter in the same column indicates no significant difference based on HSD test (5%). Above the average figure has been transformed using arcsin. J. Trop. Plant Prot.

1 (1): 46-51 (2015) 49 The Disease Intensity of CMV on Tomato Plants The data showed that the disease intensity of CMV on tomato plant without application of M. jalapa leaf extract was higher (50,2 % in average) than other tomato plants treated by M. jalapa leaf extract application. The data also showed that the higher the concentration of M. jalapa leaf extract application the lower the disease intensity of CMV.

The disease intensity of CMV on tomato plants treated with *M. jalapa* leaf extract were ranging at 31-45% (Table 2). The above results indicated that leaf extracts of *M. jalapa* decreased the disease intensity of CMV infected on tomato plants. In accordance with previous result that application of *M.*

*jalapa* leaf extract could delay the incubation period of CMV disease, it can be suggested that the application of *M. jalapa* leaf extract showed to inhibit the development of CMV disease symptom. Dean and Kuc (1986) and Suganda et al. (2002) stated that the induction of resistance from external treatment did not make the plants become immune or not affected at all, but simply increase the degree of resistance and inhibit the progression of the disease. The ability of *M.*

*jalapa* leaf extract in inducing plant defense against plant disease is suspected to be an active compound contained in *M. jalapa* which induces defense related genes such as pathogenesis-related (PR) genes which encode pathogenesis-related proteins i.e peroxidase and catalase enzyme present in tomato plants (Murphy et al., 2001).

In this study, we could not conclude whether the inhibition of CMV disease intensity was related to the inhibition of virus titer in plants, since we did not measure the CMV virus titer in tomato plants. According to Nicks (1993) the extent of damage or symptoms in plants is not always positively correlated with the level of concentration of viruses in plants.

**Peroxidase Enzyme Activity** The data showed that the activity of peroxidase enzyme activity was higher on tomato plants treated with leaf extract of *M. jalapa* compared to that of without application of *M. jalapa* leaf extract (control) particularly at 1 and 14 days after application (Figure 2). Van Loon et al. (1994) stated that peroxidase, a member of PR-protein from the group of PR-9, was accumulated at the time of plants infected by plant pathogen including virus.

Gupta et al. (1990) stated that the resistant plants against plant disease tend to exhibit higher peroxidase activity compared to the susceptible plants. Zhou et al. (1992) also stated that increased expression of peroxidase activity caused by infection of plant pathogenic virus was correlated with the level of resistance to the virus. **Catalase Enzyme Activity** The data showed that the catalase enzyme activity was elevated on tomato plants treated with *M.*

*jalapa* leaf extract particularly at 1 and 14 days after application. The average of the catalase enzyme activity in tomato plants was induced 1.3 - 1.5 times higher than in the untreated (control) tomato plants. Figure 2. **Peroxidase Enzyme Activity In Tomato Plant**

Leaves J. Trop. Plant Prot. 1 (1): 46-51 (2015) 50 Figure 3.

Catalase **Enzyme Activity In Tomato Plant** Leaves Catalase enzyme has been widely known essential to destroy H<sub>2</sub>O<sub>2</sub> formed in the peroxisomes via the oxidation reaction. H<sub>2</sub>O<sub>2</sub> is known as oxidative agent that **involved in the development of** necrotic lesion as part of disease symptom development. Therefore, the increase of catalase activity will decrease the development of symptom caused by plant pathogen.

Increased activity of catalase in tomato plants induced by leaf extract of *M. jalapa* suggested that the antioxidant induction **effect of *M. jalapa* leaf extract** help the defense reaction against CMV infection. CONCLUSION Based on the results of the study, **it can be concluded that the application of *M. jalapa* leaf extract could inhibit the development of CMV disease in tomato plants. The higher the concentration of *M.***

***jalapa* leaf extract applied on tomato plant, the lower disease incidence as well as the longer incubation period of CMV disease** disease in tomato plants. The **applications *M. jalapa* leaf extract elevated the activity of peroxidase and catalase enzymes. Thus, the inhibition of CMV disease development in tomato plant by application of *M.***

***jalapa* leaf extract may related to the induction of peroxidase and catalase activity in tomato plants.** REFERENCES Abadi, A. L. 2003. Ilmu Penyakit Tumbuhan Jilid 3. Bayumedia. Malang. 135 Hlm. Agrios, G. N. 2005. Ilmu Penyakit Tumbuhan. Edisi Kelima. Elsevier Academic Press. USA. 921 Hlm. Crescenzi, A., **Barbarossa L, Cillo F, Di Franco A, Vovlas N,** & Gallitelli D. 1993.

**Role of Cucumber mosaic virus and its satellite RNA in the etiology of tomato fruit necrosis in Italy. Arch Virol 131: 321-333. Dean, R. and J. Kuc. 1986. Induced systemic protection in cucumber : Time of production and movement of the signal. Phytopathology 61:1110-1112. Gallitelli, D. 1998. Present status of controlling cucumber mosaic virus. In : Khetarpal RK, Kagonezawa H, Hadidi A, Ed. Control of Plant Virus Diseases. APS Press, St. Paul, MN, USA.**

Gupta, S.K., P.P. Gupta, T.P. Yadava, and C.D. Kaushik. 1990. Metabolic changes in mustard due to *Alternaria* leaf blight. **Indian Phytopathol. 43(1): 64-69. Hersanti, Ling. F., dan Zulkarnain. I. 2004. Pengaruh Ekstrak Beberapa Tumbuhan dalam Menginduksi Ketahanan Sistemik Tanaman Cabai Merah ( *Capsicum annum* L. ) terhadap Cucumber Mosaic Virus (CMV). Disertasi . Sekolah Pascasarjana. Universitas Padjadjaran. 115 Hlm. J. Trop. Plant Prot.**

1 (1): 46-51 (2015) 51 Kartapradja, **R. dan D. Djuariah, 1992. Pengaruh tingkat**

kematangan buah tomat terhadap daya kecambah, pertumbuhan dan hasil tomat. Buletin Penelitian Hortikultura Vol XXIV/2. Lecoq, H., G. Wisler dan M. Pitrat. 1998. Cucurbit Viruses : The classics and The Emerging. INRA, station de Pathologie Vegetable, Domaine Saint Maurice, BP 94, 84143 Montfavet cedex. France Luck, H., 1974, Methods of Enzymatic Analysis; Bergmeyer, H. U.;

Gawehn, K., Eds.; Academic: New York, pp 885894 Murphy, A. 2001. Signal transduction in resistance to plant viruses. Euro. J.Plant Pathol. 107: 121-128. Phytopathology 66:204-208. Purwati, E. dan A. Asga, 1990. Seleksi varietas tomat untuk perbaikan kualitas. Buletin Penelitian Hortikultura Vol XX/1. Reddy, K.J., L. Wang. And S. P Gloss. 1995. Solubility and mobility of copper, zinc and lead in acidic environments. Plant and Soil. 171, 53-58. Suganda, T. 1999.

Imunisasi Tanaman : Perspektif Baru untuk Melindungi Tanaman Dari Serangan Penyakit. J. Bion. 1:46-60. Van Loon, L.C., W.S. Pierpoint.W.S., Boller.T.H., and Conejero.V. 1994. Recommendations for naming plant pathogenesis-related proteins . Plant Molecular Biology Report. 12 : 245-264. Zhou, B.W., s.Y. Liu, D.Y. Chen, Q. Yu, J. Yang, and C. Wang. 1992. Peroxidase in relation to varietal resistance to virus disease in rapeseed ( Brassica napus ). (Abstract). Oil Crops of China 2 : 52-54.

#### INTERNET SOURCES:

-----  
10% - [http://jtpp.ub.ac.id/index.php/jtpp/oai?verb=ListRecords&metadataPrefix=oai\\_dc](http://jtpp.ub.ac.id/index.php/jtpp/oai?verb=ListRecords&metadataPrefix=oai_dc)  
1% - <http://agrivita.ub.ac.id/index.php/agrivita/about/contact>  
<1% - <https://www.science.gov/topicpages/l/lankan+cassava+mosaic.html>  
3% - <http://jtpp.ub.ac.id/index.php/jtpp/article/view/116>  
1% - <https://www.bbc.com/bitesize/guides/z2gccdm/revision/3>  
<1% - <https://www.onlinejournal.in/IJIRV3I9/016.pdf>  
<1% - <https://patents.google.com/patent/US7432420B2/en>  
<1% - <https://www.sciencedirect.com/science/article/pii/B9780123943149000130>  
<1% - [https://espace.library.uq.edu.au/view/UQ:369281/UQ369281\\_OA.pdf](https://espace.library.uq.edu.au/view/UQ:369281/UQ369281_OA.pdf)  
<1% -  
<https://www.dovepress.com/challenges-and-prospects-for-the-control-of-foot-and-mouth-disease-an-peer-reviewed-fulltext-article-VMRR>  
<1% - <https://www.sciencedirect.com/science/article/pii/B978012802147700053X>  
<1% -  
[https://mafiadoc.com/naveen-kumar-arora-editor\\_599951711723dd96bd931735.html](https://mafiadoc.com/naveen-kumar-arora-editor_599951711723dd96bd931735.html)  
<1% - <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4354729/>

<1% - <http://scialert.net/fulltext/?doi=ijar.2010.954.983>  
<1% - <http://scialert.net/fulltext/?doi=ppj.2009.42.52>  
<1% - <http://protein.bio.msu.ru/biokhimiya/contents/v71/full/71S10044.html>  
<1% - <https://echa.europa.eu/registration-dossier/-/registered-dossier/25188/7/7/1>  
<1% - [https://www.researchgate.net/profile/Dennis\\_Gatahi](https://www.researchgate.net/profile/Dennis_Gatahi)  
1% - [https://mafiadoc.com/fapa-congress-2016\\_5a78b8451723dd82dd85f0d1.html](https://mafiadoc.com/fapa-congress-2016_5a78b8451723dd82dd85f0d1.html)  
<1% - <http://www.mdpi.com/2073-4441/8/4/127/htm>  
<1% - <https://quizlet.com/54374893/hz-flash-cards/>  
<1% - <http://users.ba.cnr.it/VitoLinsalata/artichoke2000/abstract.html>  
<1% - <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2515528/>  
<1% - [https://www.researchgate.net/profile/Olusegun\\_Balogun2](https://www.researchgate.net/profile/Olusegun_Balogun2)  
<1% -  
[https://www.researchgate.net/publication/261597447\\_Correlative\\_evidence\\_for\\_peroxidase\\_involvement\\_in\\_disease\\_resistance\\_against\\_Alternaria\\_leaf\\_blight\\_of\\_tomato](https://www.researchgate.net/publication/261597447_Correlative_evidence_for_peroxidase_involvement_in_disease_resistance_against_Alternaria_leaf_blight_of_tomato)  
<1% - <http://www.biologydiscussion.com/viruses/study-notes-on-viruses-biology/34217>  
<1% -  
<https://www.apsnet.org/edcenter/intropp/lessons/fungi/Oomycetes/Pages/LateBlight.aspx>  
<1% - <https://www.scribd.com/document/49559423/Farmer-s-Manual>  
<1% -  
[https://www.researchgate.net/publication/276399907\\_Infeksi\\_Bean\\_common\\_mosaic\\_virus\\_pada\\_Umur\\_Tanaman\\_Kacang\\_Panjang\\_yang\\_Berbeda](https://www.researchgate.net/publication/276399907_Infeksi_Bean_common_mosaic_virus_pada_Umur_Tanaman_Kacang_Panjang_yang_Berbeda)  
1% -  
<https://www.coursehero.com/file/p3r6akc/1-Function-shown-in-Figure-is-a-b-c-d-2-The-simplified-expression-of-full-adder/>  
<1% - <http://cuke.hort.ncsu.edu/cgc/conferences/cuc96dise.html>  
<1% - <https://link.springer.com/article/10.1007/s10658-008-9376-0>  
<1% -  
[https://www.researchgate.net/publication/234040848\\_Molecular\\_and\\_Immunological\\_Characterization\\_of\\_Gluten\\_Proteins\\_Isolated\\_from\\_Oat\\_Cultivars\\_That\\_Differ\\_in\\_Toxicity\\_for\\_Celiac\\_Disease](https://www.researchgate.net/publication/234040848_Molecular_and_Immunological_Characterization_of_Gluten_Proteins_Isolated_from_Oat_Cultivars_That_Differ_in_Toxicity_for_Celiac_Disease)  
<1% -  
[https://www.researchgate.net/publication/287681481\\_Growth\\_and\\_yield\\_responses\\_of\\_strawberry\\_plants\\_to\\_chitosan\\_application](https://www.researchgate.net/publication/287681481_Growth_and_yield_responses_of_strawberry_plants_to_chitosan_application)  
<1% - [https://issuu.com/alkasingh8/docs/ajbs\\_\\_11\\_\\_1\\_](https://issuu.com/alkasingh8/docs/ajbs__11__1_)  
<1% - <https://epdf.tips/natural-resistance-mechanisms-of-plants-to-viruses.html>  
<1% - [https://en.wikipedia.org/wiki/Management\\_of\\_HIV/AIDS](https://en.wikipedia.org/wiki/Management_of_HIV/AIDS)  
<1% -  
[https://www.researchgate.net/publication/287813430\\_The\\_Pathogenesis\\_Related\\_Class\\_10\\_proteins\\_in\\_Plant\\_Defense\\_against\\_Biotic\\_and\\_Abiotic\\_Stresses](https://www.researchgate.net/publication/287813430_The_Pathogenesis_Related_Class_10_proteins_in_Plant_Defense_against_Biotic_and_Abiotic_Stresses)

<1% - <http://europepmc.org/articles/PMC4178825>  
<1% - [https://link.springer.com/chapter/10.1007%2F978-1-4614-4806-8\\_10](https://link.springer.com/chapter/10.1007%2F978-1-4614-4806-8_10)  
<1% - [http://file.scirp.org/Html/2-2601059\\_42434.htm](http://file.scirp.org/Html/2-2601059_42434.htm)  
<1% -  
[https://www.researchgate.net/publication/303742378\\_Spraying\\_of\\_leaf-colonizing\\_Bacillus\\_amyloliquefaciens\\_protects\\_pepper\\_from\\_Cucumber\\_mosaic\\_virus](https://www.researchgate.net/publication/303742378_Spraying_of_leaf-colonizing_Bacillus_amyloliquefaciens_protects_pepper_from_Cucumber_mosaic_virus)  
<1% - <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC35165/>  
<1% -  
[https://www.researchgate.net/publication/289528727\\_Effect\\_of\\_Selenium\\_on\\_Elemental\\_Concentration\\_and\\_Antioxidant\\_Enzymatic\\_Activity\\_of\\_Tomato\\_Plants](https://www.researchgate.net/publication/289528727_Effect_of_Selenium_on_Elemental_Concentration_and_Antioxidant_Enzymatic_Activity_of_Tomato_Plants)  
<1% - <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4500917/>  
<1% - [https://en.wikipedia.org/wiki/Oxidative\\_stress](https://en.wikipedia.org/wiki/Oxidative_stress)  
<1% - <https://link.springer.com/article/10.1007/s10658-017-1332-4>  
1% - [https://link.springer.com/chapter/10.1007/978-3-662-03506-1\\_12](https://link.springer.com/chapter/10.1007/978-3-662-03506-1_12)  
<1% - <https://rd.springer.com/content/pdf/10.1023%2FA%3A1009615807781.pdf>  
<1% - <https://jurnal.ugm.ac.id/jpti/article/view/12104>  
<1% - <https://fr.scribd.com/doc/89511734/Sumber-Buku>  
1% - <http://luqmanmaniabgt.blogspot.com/2012/10/makalah-kultur-tomat.html>  
1% - <https://www.tandfonline.com/doi/full/10.1080/00103624.2018.1435682>