

Functional Characteristics

by - -

Submission date: 01-Feb-2024 12:17PM (UTC+0500)

Submission ID: 2283584984

File name: anam_2023-Functional_Characteristics_of_Trash_Fish_in.pdf (464.9K)

Word count: 5276

Character count: 27188

Functional Characteristics of Trash Fish in Lamongan Regency, East Java, Indonesia

Choirul Anam^{1,*}, Noor Harini², Damat Damat², Roy Hendroko Setyobudi², Ida Ekawati³, Taavi Liblik⁴, Endang Dwi Purbajanti⁵, Hendrikus Bernedektus⁶, Loli Melani Souripet⁷, Ahmad Fauzi², Afrida Rizka Farzana⁸, Rusli Tonda⁹, Iswahyudi Iswahyudi¹⁰, Ana Amiroh¹, Mariyatul Qibtiyah¹, Dian Eka Kusumawati¹, Istiqomah Istiqomah¹, and Emmy Hamidah¹

¹Universitas Islam Darul 'Ulum, Lamongan 62253, East Java, Indonesia

²University of Muhammadiyah Malang, Malang 65144, East Java, Indonesia

³Universitas of Wiraraja, Sumenep 69451, East Java, Indonesia

⁴Tallinn University of Technology, 12616 Tallinn, Estonia

⁵University of Diponegoro, Semarang 50275, Central Java, Indonesia

⁶Department of Agriculture and Food of Papua Province, Jayapura 99112, Indonesia

⁷Department of Food Crops, Horticulture, and Plantations, West Papua Province, Mamberi 98312, Indonesia

⁸IPB University, Bogor 16680, West Java, Indonesia

⁹University of Tribhuwana Tungadewi, Malang 65144, East Java, Indonesia

¹⁰Universitas Islam Madura, Pamekasan 69317, East Java, Indonesia

Abstract. Trash fish has several weaknesses, such as having little meat, and being highly perishable. The purpose of this study was to determine the functional characteristics of trash fish through fillet preparation techniques and types of fish. This study used a completely randomized design method. The first factor is the type of fish consisting of Orangefin ponyfish (*Leiognathus bindus* Valenciennes, 1835), Chacunda gizzard-shad (*Anodontostoma chacunda* Hamilton, 1822) and Sardine (*Sardinella Fimbriata* Valenciennes, 1847). The second factor was the type of preparation technique which consisted of mechanical, blanching, 1 % acid immersion and 1 % papain enzyme immersion. The data obtained were then analyzed descriptively from the preparation technique for each observation parameter and presented in tabular form and plotted in graphical form. The results of this study indicate that Orangefin ponyfish, Chacunda gizzard-shad and Sardine fish have a range of functional properties of the three types of fish, namely: foaming power (17.68 % to 61.87 %), foam stability (50 % to 57.14 %), emulsifying power (3.31 % to 4.29 %), emulsion stability (1.91 % to 3.37 %), WHC (33.9 % to 46.64 %), and OHC (24.75 % to 29.57 %).

Keywords: Fish fillet functional properties, preparation technique, trash fish, waste utilization.

* Corresponding author: choirulanam@unisda.ac.id

1 Introduction

Lamongan regency is known as one of the fisheries areas or minapolitan in East Java, Indonesia [1, 2]. Fish that are not included in the main purpose of fishing are called trash fish and are commonly used as a mixed ingredient in the production of animal feed [3] or processed into salted fish and sometimes just thrown away resulting in a rotten smell during the main harvest season. Several researchers have recommended trash fish to be a product diversification as an effort to recycle or waste utilization [4–8]

Trash fish in Lamongan has the potential to be made into high-value economic products, namely Orangefin ponyfish (*Leiognathus bindus* Valenciennes, 1835), Chacunda gizzard-shad (*Anodontostoma chacunda* Hamilton, 1822) and Sardine (*Sardinella Fimbriata* Valenciennes, 1847) [9]. Trash fish has a small and varied morphology, is easily damaged, has tight spines, and has little flesh, making it difficult to separate the fish flesh from the bones and skin. Trash fish is considered a marine fisheries waste [10].

Animal food products from fish meat make a very important contribution as a source of protein. Protein with its functional properties can make food ingredients very attractive and encourage the emergence of flavors, textures and other qualities desired by consumers [11]. Fresh fish can be converted into semi-finished raw materials, namely in the form of fillets, surimi and so on.

To obtain fillets with good physical, chemical and functional characteristics, various preparation techniques are necessary. Kuniran fish (*Upeneus sulphureus* Cuvier, 1829) preparation technique by slashing has a high yield value (67.50 %), and a faster separation process time (11.35 kg h⁻¹) [12]. The preparation technique for trash fish fillets with 1 % enzyme dissolution has better physical and chemical characteristics than the mechanical preparation techniques, blanching, or with 1 % acid dissolution [13]. While functional properties: Suglir fish protein concentrate (*Elagatis bipinnulatus* Quoy & Gaimard, 1825) has oil absorption (2.48 g g⁻¹) and water absorption (2.02 mL g⁻¹) [14], in Wader fish fillets (*Rashora jacobsoni* Weber & de Beaufort, 1916) has foaming power (211.03 %), foam stability (19.17 %), emulsifying power (2.42 %), emulsion stability (0.55 %) [15], in cowpea protein isolate (*Vigna unguiculata* (L.) Walp.) has foaming power (68 mL g⁻¹), foam stability (8 %), OHC (84.89 %), WHC (136.61 %) [16]. This study aims to obtain the functional properties of Indonesian Lamongan trash fish fillets.

2 Methods

2.1 Materials

The main ingredients of fresh trash fish are Orangefin ponyfish, Chacunda gizzard-shad and Sardine with an average weight of 50 g tail⁻¹ to 100 g tail⁻¹ purchased from the fish mother market in Lamongan Regency, East Java, Indonesia. The fish was immediately stored in the refrigerator and transported to the Department of Agroindustry Technology, the University of Jember for 52. Once you arrive, the fish is stored in the freezer. Before analysis, the fish is immediately thawed at room temperature, washed, filled, and chopped to a uniform using a chopper. The ingredients used are young papaya fruit, lime juice, cooking oil, CH₃COOH and other chemicals (analytical class) are TCA 5 %, sodium hydroxide, H₂SO₄, selenium, boric acid 3 %, SDS 0.1 %, and petroleum benzene [15].

2.2 Preparation of trash fish fillets

Preparation fish samples to eliminate fishy odor by giving lime as much as 15 % (b/v) of the weight of the fish for 10 min. Removing fishy odor in the trash done after washing the fish with clean water. The three types of marine fish have carried out preparation techniques in four ways, namely mechanical preparation techniques with the Palmeira method in fish [17], blanching with the Nguyen method on asparagus [18] and chemically with the Moniharapon method in fish [19], and enzymatic with the Ma method on softening meat [20]. The preparation technique performed on fish are given the same time limit of 5 min for blanching treatment and 30 min for chemical and enzyme treatments. Preparation fish mechanically is done by way of fillets directly on fresh fish. In this process, the three types of fish are carried out early washing using running water, while the other fish are temporarily accommodated in a freezer. Fish have been washed discarded scales and stomach contents, then the fish is carried out the filleted process from the tip of the tail, splitting the back to the direction of the head. Fish that have been cleaned and detached from thorns are weighed and recorded in weight stored in a freezer.




Fig. 1. Flowchart of trash fish preparation technique [21].

In the technique of fish blanching, all three types of fish are done by way of fish cleaned alternately using running water. After washing, the fish is boiled at a temperature of 100 °C for 5 min. The fish is lifted and cooled in the open air. After chold, the three types of fish have carried out the removal of thorns, and the fillet process is carried out. Fish that are clean and detached from thorns are weighed and recorded in weight the stored in the freezer. Chemical and enzymatic preparation techniques are carried out by soaking fish. Chemically, fish is soaked with acetic acid solution at a concentration of 1 %. Enzymatically, fish are soaked with a solution of papain enzymes at concentration of 1 %. This immersion is done for 30 min, after which the fillet process is then weighed and recorded the results. The process of filleting trash fish can be seen in Figure 1.

Papain enzymes are made by tapping the fruit skin of papaya plants that have been aged 2 mo to 3 mo. Tapping is done by sticking a sad tool (knife) on the fruit's skin from the base to the end of the fruit. The depth of the nick is between one mm to mm. Papaya sap that comes out is accommodated in a container. Papaya sap mixed with buffer phosphate 0.2 M p14 7 (ratio 1:1). The mixture is centrifuged at 8 000 rad s⁻¹ at 4 °C for 10 min (1 rad s⁻¹ = 1/60 Hz). Filtrate is a papain enzyme. The manufacture of papain enzyme follows the method of Urgessa [22] with a little modification.

The yield of the preparation technique on each treatment will be analyzed for physical characteristics and 15 chemical. Chemical and enzymatic treatment will be taken at the concentration with the highest yield and the fastest time based on the fillet type of trash fish.

2.3 Functional properties

Analysis for 10 foaming capacity and foam stability [16], emulsion capacity and emulsion stability [23], water holding capacity [24], and oil holding capacity [16].

2.4 Data analysis

The research design used was experimental laboratory. First, the test data is calculated using Microsoft Excel and presented in table form. Next, it is explored graph form, then analyzed with the ANOVA - One Way (minitab 17) test. After that, Tukey test ($P \leq 0.05$) if there is a significant difference [25, 26]

3 Results and discussion

3.1 Foam strength and stability

The results of variance showed that variations in preparation techniques and fish species had a significant effect on foaming capacity and there was an interaction between the two factors. 16 foaming ability of Sardine, Chacunda gizzard-shad and Orangefin ponyfish fish fillets can be seen in Figure 2. Meanwhile, the results of the foam stability analysis showed that variations in preparation techniques and types of fish had a significant effect and there were interactions between the two factors. The foam stability of Sardine, Chacunda gizzard-shad and Orangefin ponyfish fish fillets is shown in Figure 3.




Fig. 2. Fish fillet foaming capacity.

Figure 2 shows the foaming power contained in Sardine, Chacunda gizzard-shad and Oranegin ponyfish fish fillets with mechanical treatment (17.68 %, 21.82 %, and 30 %). The foaming capacity of Sardine, Chacunda gizzard-shad and Oranegin ponyfish fish fillets in the blanching treatment (10 %, 10 %, and 0 %). The foaming ability of Sardine, Chacunda gizzard-shad and Oranegin ponyfish fish fillets in 1 % acetic acid solution immersion (24.09 %, 26.51 %, and 18.28 %). Sardine, Chacunda gizzard-shad and Oranegin ponyfish fulllet foaming ability in 1 % papain enzyme solution immersion treatment (18.79 %, 61.87 %, and 17.68 %). The highest foaming power was found in Chacunda gizzard-shad fish treated with 1 % enzyme, which was 61.87 %. This value was significantly different compared to Chacunda gizzard-shade fish with 1 % acid treatment, which was 26.51 %.

The results of the analysis above show that the foaming power of Sardine, Chacunda gizzard-shad and Oranegin ponyfish fish in the mechanical treatment, blanching, soaking in acid solution and immersion in papain enzyme solution is lower when compared to the foaming capacity of Wader fish fillets, which is equal to 211.03 % [15]. The foaming power value will be lower after the processing treatment. This is because the treatment during processing can dissolve water-soluble sarcoplasmic proteins, causing changes that lead to a decrease in foaming power. Sarcoplasmic protein is a reflection of the relationship between protein solubility and foam formation [27]. This is in accordance with the statement of Afrianto et al. [28], blanching by boiling has weaknesses, including being able to remove nutrients in water-soluble fish, in this case the foaming ability of the three types of blanching treated fish has a low value.




Fig. 3. Foam stability of fish fillet.

Figure 3 shows the stability of the foam contained in Sardine, Chacunda gizzard-shad and Orangefin ponyfish fish fillets with mechanical treatment (0 %, 44.44 %, and 33.33 %). The foam stability of the blanching treatment of all types of fish was 0 %. The foam stability of Sardine, Chacunda gizzard-shad and Orangefin ponyfish fish fillets in the 1 % acetic acid solution immersion treatment (38.89 %, 33.33 %, and 50 %). Foam stability of Sardine, Chacunda gizzard-shad and Orangefin ponyfish fish fillets in 1 % papain enzyme solution immersion (50 %, 57.14 %, and 50 %). The highest foam stability was found in Chacunda gizzard-shad fish treated with 1 % enzyme, which was 57.14 %. This value was significantly different from the stability of the foam of Chacunda gizzard-shad fish with mechanical treatment, which was 44.44 %.

The results of the analysis above show that the foam stability of Sardine, Chacunda gizzard-shad and Orangefin ponyfish fish in mechanical treatment, soaking in acid solution and soaking in papain enzyme solution is higher when compared to the foam stability of Wader fish which is 19.17 %, and Bader fish (*Puintius javanicus* Bleeker, 1855) namely 18.38 % [15]. While fish with mechanical treatment and all types of fish with blanching treatment have a lower value.

The low value of foam stability is due to the foam that is formed large and breaks easily. Foam strength and stability increase with increasing protein concentration, where increasing protein concentration can increase viscosity and form many layers of protein cohesion on the surface. Foaming capacity is influenced by intramolecular and flexibility of protein molecules [29].

3.2 Emulsion capacity and stability

The emulsion capacity of Sardine, Chacunda gizzard-shad and Orangefin ponyfish fish fillets ranged from 2.64 % to 5.94 %. The results of the analysis of variance showed that there was an interaction between the preparation technique and the type of fish on the emulsifying

power of the fish fillets. The emulsion power of Sardine, Chacunda gizzard-shad and Orangefin ponyfish fish fillets can be seen in Figure 4, while the emulsion stability of Sardine, Chacunda gizzard-shad and Orangefin ponyfish fish fillets ranges from 1.56 % to 3.37 %. The results of the emulsion stability analysis of variance showed that there was no interaction between the various preparation techniques and the types of fish. The emulsion stability of Sardine, Chacunda gizzard-shad and Orangefin ponyfish fish fillets can be seen in Figure 5.




Fig. 4. Emulsion capacity of fish fillets.

Figure 4 shows the emulsifying power contained in Sardine, Chacunda gizzard-shad and Orangefin ponyfish fish fillets with mechanical treatment (3.82 %, 3.07 %, and 4.62 %). Emulsion power of Sardine, Chacunda gizzard-shad and Orangefin ponyfish fish fillets in the blanching treatment (5.67 %, 5.94 %, and 5.21 %). Emulsion power of Sardine, Chacunda gizzard-shad and Orangefin ponyfish fish fillets in 1 % vinegar solution (2.64 %, 3.36 %, and 3.67 %) immersion treatment. Meanwhile, the emulsion power of Sardine, Chacunda gizzard-shad and Orangefin ponyfish fish fillets was treated with 1 % papain enzyme solution immersion (3.31 %, 4.29 %, and 4.14 %). The highest emulsion power was found in the blanching treatment of Chacunda gizzard-shad fish, which was 5.94 %. This value was significantly different compared to the emulsion power of Chacunda gizzard-shad fish with 1 % enzyme treatment, which was 4.29 %. The three types of fish had the ability to form good emulsions in the blanching treatment, while the other types of fish had almost the same emulsion power (2.64 % to 4.29 %). The emulsion power of all three types of fish treatment was still higher when compared to Wader fish which had an emulsion power of 2.42 % [15]. Emulsion depends on the high absorption capacity of oil-water ($w\ o^{-1}$), and proteins with a high amount of hydrophobicity will be adsorbed on the interfacial oil-water ($w\ o^{-1}$) interface. Protein will lower the surface tension between the phases and form an emulsion [30]. This can determine the formation of oil and water emulsions as well as in the stabilization process.




Fig. 5. Fish fillet emulsion stability.

Figure 5 shows the stability of the emulsion found in Sardine, Chacunda gizzard-shad and Orangefin ponyfish fish fillets with mechanical treatment (1.56 %, 2.28 %, and 1.82 %). Emulsion stability of Sardine, Chacunda gizzard-shad and Orangefin ponyfish fish fillets in the blanching treatment (2.32 %, 1.81 %, and 1.89 %). Emulsion stability of Sardine, Chacunda gizzard-shad and Orangefin ponyfish fish fillets in 1 % vinegar solution immersion (1.98 %, 2.14 %, and 1.63 %). Stability of Sardine, Chacunda gizzard-shad and Orangefin ponyfish fish fillet emulsion in 1 % papain enzyme solution immersion treatment (1.91 %, 1.93 %, and 3.37 %). The highest emulsion stability was found in Orangefin ponyfish fish fillet with 1 % enzyme treatment, namely 3.37 %. This value was significantly different from the stability of the blanching treatment of Orangefin ponyfish fish emulsion, which was 1.89 %. The three types of fish have higher emulsion stability when compared to Wader fish which has an emulsion stability of 0.55 % [15]. The stability of the emulsion depends on the thickness and thickness of the membrane or protein film absorbed by the interface between the oil and water phases. Emulsification properties are determined by the quality of the protein, not the quantity or amount of protein [31]. Emulsion stability depends on the interpartial strength of the ingredients in maintaining the hydrophobic interaction between oil and protein [32]. Emulsion stability is affected by emulsification conditions in proteins, protein sources and concentrations, pH, ionic strength (salt type and concentration) and viscosity in the food system [33].

3.3 Water Holding Capacity (WHC)

The calculation results obtained WHC values for Sardine, Chacunda gizzard-shad and Orangefin ponyfish fish fillets in the mechanical treatment (42.72 %, 34.697 %, and 27.64 %). WHC values in the blanching treatment of Sardine, Chacunda gizzard-shad and Orangefin ponyfish fish fillets (78.9 %, 112.39 %, and 43.24 %). WHC values of Sardine, Chacunda gizzard-shad and Orangefin ponyfish fish fillets in 1 % acid solution immersion treatment (35.96 %, 17.98 %, and 44.44 %). WHC values in 1 % enzyme solution immersion

treatment of Sardine, Chacunda gizzard-shad and Orangefin ponyfish fish fillets (33.9 %, 43.11 %, and 46.64 %). The results of variance showed that variations in preparation techniques and fish species had a significant effect and there was an interaction between the two parameters (Figure 6).




Fig. 6. WHC value of fish fillets.

Figure 6 shows the largest WHC value is in Chacunda gizzard-shad fish fillets with blanching treatment (112.39 %). This value was significantly different when compared to Chacunda gizzard-shad fish fillets treated with 1 % enzyme (43.11 %). This could be due to the bond in the fish fillet tissue which has changed due to the influence of high temperatures, making it easy to absorb water. In addition, the WHC value is influenced by the protein content of the meat. Chacunda gizzard-shad fish has a good protein content (8.89 % to 10.52 %) [13] so it can hold water in the fish meat. Water absorption is strongly influenced by protein concentration, ionic strength, temperature and other components such as hydrophilic polysaccharides, fats, salts, and is also influenced by the duration of heating and storage conditions. [13] higher the protein concentration, the better the water absorption. As the protein content of the meat increases, the WHC of the meat will increase due to the protein's ability to chemically bind water [34]. The ability to absorb or bind water is needed, especially in one of the most important food applications, for example in the manufacture of sausages.

Mechanical treatment, immersion in vinegar and enzyme solutions has a relatively low water absorption capacity. This can be caused by the low protein content in fish fillets and due to a decrease in pH. The low WHC value is caused by changes in the ions bound by the meat protein due to the large amount of lactic acid that has accumulated as a result of which many myofibril proteins are damaged. In addition, the hydrolysis of meat protein by enzymes causes the volume of muscle fibers to expand. This damage causes the loss of the protein's ability to bind water [35].

3.4 Oil Holding Capacity (OHC)

The calculation results obtained OHC values for Sardine, Chacunda gizzard-shad and Orange-fin ponyfish fish fillets in the mechanical treatment (34.55 %, 23.25 %, and 13.33 %). The OHC value of Sardine, Chacunda gizzard-shad and Orange-fin ponyfish fish fillets in the blanching treatment (59.14 %, 71.52 %, and 55.63 %). OHC values of Sardine, Chacunda gizzard-shad and Orange-fin ponyfish fish fillets in 1 % acid solution immersion treatment (8.64 %, 21.94 %, and 12.58 %). OHC values of Sardine, Chacunda gizzard-shad and Orange-fin ponyfish fish fillets in 1 % enzyme solution immersion treatment (29.57 %, 24.75 %, and 26.49 %). The results of the variance indicated that there was an interaction between various preparation techniques and fish species (Figure 7).




Fig. 7. OHC value of fish fillets.

Figure 7 shows the largest OHC value was found in Chacunda gizzard-shad fish fillets with the blanching treatment (71.52 %). This value was significantly different from the OHC Juwi fish with 1 % enzyme treatment (24.75 %). This could be due to the protein content of Chacunda gizzard-shad fish in the blanching treatment having the highest protein content (10.59 %) [13] compared to other types of fish, so that Chacunda gizzard-shad fish have good oil absorption abilities. Oil absorption occurs because the oil is physically trapped in the protein. Protein structure is a factor that greatly determines the ability to absorb oil. Structures that are more lipophilic contain more non-polar protein branches and will contribute to increased oil absorption [36].

4 Conclusion

Sardine, Chacunda gizzard-shad and Orange-fin ponyfish fish have functional properties, namely: foaming power (17.68 % to 61.87 %), foam stability (50 % to 57.14 %), emulsifying power (3.31 % to 4.29 %), emulsion stability (1.91 % to 3.37 %), WHC (33.9 % to 46.64 %), and OHC (24.75 % to 29.57 %).

References

1. D.R. Raissa, R.P. Setiawan, D. Rahmawati, *Procedia-Social Behav. Sci.*, **135**: 167–171 (2014) <https://doi.org/10.1016/j.sbspro.2014.07.342>
2. M.S.A. Ningsih, P. Prayogo, A.M. Sahidu, *IOP Conf. Ser.: Earth Environ. Sci.*, **441**,012034: 1–8 (2020) <https://iopscience.iop.org/article/10.1088/1755-1315/441/1/012034>
3. M.A. Nugraha, R. Rozi, *IOP Conf. Ser.: Earth Environ. Sci.*, **441**,012069: 1–5 (2020) <https://iopscience.iop.org/article/10.1088/1755-1315/441/1/012069>
4. V. Rudovica, A. Rotter, S.P. Gaudêncio, L. Novoveská, F. Novoveská, L.K. Akslen-Hoel, et al., *Front. Mar. Sci.*, **8**,723333: 1–39 (2021) <https://doi.org/10.3389/fmars.2021.723333>
5. H. Susanto, R.H. Setyobudi, S.M. Nur, E. Yandri, A.S. Uyun, A. Yaro, et al., *E3S Web Conf.*, **190**,00014: 1–10 (2020) <https://doi.org/10.1051/e3sconf/202019000014>
6. C. Anam, M.F.M. Atoum, N. Harini, D. Damat, R.H. Setyobudi, A. Wahyudi, et al., *Jordan J. Biol. Sci.*, **16**,2: 267–277 (2023) <https://doi.org/10.54319/jjbs/160211>
7. K. Abdullah, A.S. Uyun, R. Soegeng, E. Suherman, H. Susanto, R.H. Setyobudi, et al., *E3S Web Conf.*, **188**00016: 1–8 (2020) <https://doi.org/10.1051/e3sconf/202018800016>
8. J. Burlakovs, Z. Vincevica-Gaile, V. Bisters, W. Hogland, M. Kriipsalu, I. Zekker, et al., *Application of Anaerobic Digestion for Biogas and Methane Production from Fresh Beach-Cast Biomass*. EAGE GET 2022 (The Hague, Netherlands, 2022) *Proceeding EAGE GET 2022*: 1–5 (2022) <https://doi.org/10.3997/2214-4609.202221028>
9. C. Anam, N. Harini, D. Damat, A. Wahyudi, Y. Witono, N. Kuswardhani, et al., *E3S Web Conf.*, **226**,00011: 1–11 (2021) <https://doi.org/10.1051/e3sconf/202122600011>
10. N.D. Rahayu, L. Sulmartiwi, G. Mahasri, M. Muntalim, B. Angwarmas, G.D. Pamenang, *IOP Conf. Ser.: Earth Environ. Sci.*, **441**,012095: 1–7 (2020) <https://iopscience.iop.org/article/10.1088/1755-1315/441/1/012095>
11. R.J.S. de Castro, M.A.F. Domingues, A. Ohara, P.K. Okuro, J.G. dos Santos, R.P. Brex'õ, et al., *Food Struct.*, **14**: 17–29 (2017) <http://dx.doi.org/10.1016/j.foostr.2017.05.004>
12. B.B. Sedayu, I.M.S. Erawan, P. Wullandari, J. Pascapanen *Bioteknologi Kelautan Perikanan*, **10**,1: 83–89 (2015) [in Bahasa Indonesia] <http://bbp4b.litbang.kkp.go.id/jurnal-jpbkp/index.php/jpbkp/article/view/247>
13. C. Anam, *Pengembangan teknologi produksi protein miofibril ikan rucah pantura Lamongan sebagai food ingredient* [Development of technology for the production of myofibril protein for trash fish in the north coast of Lamongan as a food ingredient] [Dissertation] Pascasarjana, Universitas Muhammadiyah Malang (2021) [in Bahasa Indonesia]. p.76 <https://eprints.umm.ac.id/80567/>
14. F.J. Rieuwpassa, E.J. Karimela, D.C. Lasaru. *J. Teknol. Perikanan Kelautan*, **9**,2: 177–183 (2019) [in Bahasa Indonesia] <https://journal.ipb.ac.id/index.php/jtpk/article/view/25774>
15. I. Akbariawati, *Karakteristik fisik, kimia, dan fungsional fillet ikan wader (Rasbora jacobsoni), Bader (Puntius javanicus), dan patin (Pangasius hypophthalmus) akibat dari perbedaan teknik preparasi* [Physical, chemical, and functional characteristics of wader (*Rasbora jacobsoni*), Bader (*Puntius javanicus*), and catfish (*Pangasius hypophthalmus*) fillets as a result of differences in preparation techniques] [Undergraduated Thesis] Teknologi Hasil Pertanian, Universitas Jember (2015) [in Bahasa Indonesia] <https://repository.unej.ac.id/handle/123456789/72479>
16. Y. Witono, C. Anam, H. Herlina, A.D. Pamujiati, *Int. J. Adv. Sci. Eng. Inf. Technol.*, **4**,2: 58–62 (2014) <https://doi.org/10.18517/ijaseit.4.2.377>

17. K.R. Palmeira, E.T. Mársico, M.L.G. Monteiro, M. Lemos, C.A. Conte Junior, *CyTA J. Food*, **14**,2: 227–238 (2016) <https://doi.org/10.1080/19476337.2015.1087050>
18. T.V.L. Nguyen, T.T. Vo, T.D. Lam, L.G. Bach, *Mater. Today: Proc.*, **18**,part 7: 4799–4809 (2019) <https://doi.org/10.1016/j.matpr.2019.07.468>
19. T. Moniharapon, F. Pattipeilohy, D.L. Moniharapon, R.B.D. Sormin, *IOP Conf. Ser.: Earth Environ. Sci.*, **339**,012051: 1–13 (2019) <https://iopscience.iop.org/article/10.1088/1755-1315/339/1/012051>
20. Y. Ma, Y. Yuan, X. Bi, L. Zhang, Y. Xing, Z. Che, *Food Bioprocess Technol.*, **12**: 681–693 (2019) <https://doi.org/10.1007/s11947-019-2245-3>
21. S.S. Chan, B. Roth, M. Skare, M. Hernar, F. Jessend, T. Løvdal, et al., *J. Aquac.*, **526**,735381: 1–10 (2020) <https://doi.org/10.1016/j.aquaculture.2020.735381>
22. O.E. Urgessa, D.D. Itana, T.O. Raga, *Ethiop. J. Sci. Sustain. Dev.*, **6**,2: 22–32 (2019) <https://doi.org/10.20372/ejssdastu:v6.i2.2019.92>
23. J.K. Parkington, Y.L. Xiong, S.P. Blanchard, S. Xiong, B. Wang, S. Srinivasan, et al., *J. Food Sci.* **65**,3: 428–433 (2008) <https://doi.org/10.1111/j.1365-2621.2000.tb16021.x>
24. Z. Zhang, Y. Yang, X. Tang, Y. Chen, Y. You, *Food Sci. Technol. Res.*, **21**,4: 597–605 (2015) <https://doi.org/10.3136/fstr.21.597>
25. P.G. Adinurani, *Perancangan dan Analisis Data Percobaan Agro: Manual and SPSS [Design and Analysis of Agrotrial Data: Manual and SPSS]* (Plantaxia, Yogyakarta, 2016) [in Bahasa Indonesia] <https://opac.perpusnas.go.id/DetailOpac.aspx?id=1159798#>
26. P.G. Adinurani, *Statistik Terapan Agroteknologi (disusun sesuai rencana pembelajaran semester) [Agrotechnology Applied Statistics (compiled according to the semester learning plan)]* (Deepublish, Yogyakarta, 2022) [in Bahasa Indonesia] <https://deepublishstore.com/shop/buku-statistika-terapan-3/>
27. C. Simeanu, D. Simeanu, A. Popa, A. Usturoi, D. Bodescu, M. G. Dolis, *Rev. Chim.*, **68**,5: 1063–1069 (2017) <https://doi.org/10.37358/rc.17.5.5612>
28. E. Afrianto, E. Liviawaty, O. Suhara, H. Hamdani. *J. Akuatika*, **5**,1: 45–54 (2014) [in Bahasa Indonesia] <http://jurnal.unpad.ac.id/akuatika/article/view/3704>
29. N. Phawaphuthanon, D. Yu, P. Ngamnikom, I.S. Shin, D. Chung, *Food Hydrocoll.*, **88**: 119–126 (2019) <https://doi.org/10.1016/j.foodhyd.2018.09.041>
30. M. Padiál-Domínguez, F.J. Espejo-Carpio, R. Pérez-Gálvez, A. Guadix, E.M. Guadix, *Foods*. **9**,5: 1–15 (2020) <https://doi.org/10.3390/foods9050636>
31. S. Sathivel, P.J. Bechtel, *J. Food Biochem.*, **32**,5: 557–575 (2008) <https://doi.org/10.1111/j.1745-4514.2008.00184.x>
32. J. Zhao, J. Li, J. Wang, W. Lv, *J. Agric. Food Chem.*, **60**,45: 11387–11394 (2012) <https://doi.org/10.1021/jf303439p>
33. V.S. Krstonošić, M.D. Kalić, T.R. Dapčević-Hadnađev, I.S. Lončarević, M.S. Hadnađev, *Colloids Surf. A: Physicochem. Eng. Asp.*, **602**,125045: 1–33 (2020) <https://doi.org/10.1016/j.colsurfa.2020.125045>
34. Z.H. Duan, H.Z. Liu, P. Luo, Y.P. Gu, Y.Q. Li, *Chem. Cent. J.*, **12**,30: 1–7 (2018) <https://doi.org/10.1186/s13065-018-0402-9>
35. B. Rezaharsamto, E. Subroto, *Int. J. Sci. Technol. Res.*, **8**,12: 3151–3156 (2019) <https://www.ijstr.org/final-print/dec2019/-A-Review-On-Bioactive-Peptides-Derived-From-Various-Sources-Of-Meat-And-Meat-By-products.pdf>
36. D. Ananey-Obiri, L. Matthews, M.H. Azahrani, S.A. Ibrahim, C.M. Galanakis, R. Tahergorabi, *Trends Food Sci. Technol.*, **80**: 167–174 (2018) <https://doi.org/10.1016/j.tifs.2018.08.012>

Functional Characteristics

ORIGINALITY REPORT

10%

SIMILARITY INDEX

8%

INTERNET SOURCES

9%

PUBLICATIONS

4%

STUDENT PAPERS

PRIMARY SOURCES

1	Submitted to Universitas Negeri Jakarta Student Paper	3%
2	jjbs.hu.edu.jo Internet Source	1%
3	I. Ketut Widnyana, Putu Eka Pasmidi Ariati, I. Ketut Sumantra, I. Made Wahyu Wijaya et al. "The Effect of Liquid Organic Fertilizer from Plant Waste, Livestock Waste, and Fish Waste on Growth of Marigold", E3S Web of Conferences, 2023 Publication	1%
4	Choirul Anam, Noor Harini, Damat Damat, Ahmad Wahyudi et al. "Potential Analysis of Low Economic Value Fish in Lamongan Regency, East Java, Indonesia", E3S Web of Conferences, 2021 Publication	<1%
5	www.paspk.org Internet Source	<1%
6	ejournal.unsrat.ac.id Internet Source	<1%
7	sinergiejournal.eu Internet Source	<1%
8	Chen, Liming, Warren A. Dick, and David Kost. "Circulating Fluidized Bed Combustion Product Addition to Acid Soil: Alfalfa (<i>Medicago sativa</i> L.) Composition and	<1%

Environmental Quality", Journal of
Agricultural and Food Chemistry, 2006.

Publication

9

Rusli Tonda, Manar Fayiz Mousa Atoum, Roy Hendroko Setyobudi, Lili Zalizar et al. "Food Waste Product for Overcoming Heat Stress in Broilers", E3S Web of Conferences, 2023

Publication

<1 %

10

pubs.rsc.org

Internet Source

<1 %

11

Endang Dwi Purbajanti, Didik Wisnu Widjajanto, Praptiningsih Gamawati Adinurani, Zahid Hussain, Ida Ekawati. "Production and Nutrient Value of Elephant Grass in Agroforestry Systems in Indonesia", E3S Web of Conferences, 2023

Publication

<1 %

12

eprints.umm.ac.id

Internet Source

<1 %

13

Changwei Cao, Zhichao Xiao, Huiquan Tong, Xionsong Tao, Dahai Gu, Yinglong Wu, Zhiqiang Xu, Changrong Ge. "Effect of ultrasound-assisted enzyme treatment on the quality of chicken breast meat", Food and Bioproducts Processing, 2021

Publication

<1 %

14

Ruly Sitanggang, Meiri Triani, Rasgianti Rasgianti, Eko Supriyanto, Dheka Bakti, Nur Cahyo, Maizirwan Mel. "Deposit Characteristics of Diesel Engine Power Plant Fueled by Crude Palm Oil", E3S Web of Conferences, 2023

Publication

<1 %

15

text-id.123dok.com

Internet Source

<1 %

16

vdoc.pub

Internet Source

<1 %

17

www.grafiati.com

Internet Source

<1 %

18

Roy Hendroko Setyobudi, Damat Damat, Shazma Anwar, Ahmad Fauzi et al. "Amino Acid Profiles of Coffee Cherry Flour from Different Origins: A Comparative Approach", E3S Web of Conferences, 2023

Publication

<1 %

19

digital.library.unt.edu

Internet Source

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On

Functional Characteristics

GRADEMARK REPORT

FINAL GRADE

GENERAL COMMENTS

/0

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9

PAGE 10

PAGE 11

PAGE 12
